

BOLIVIA

Cutaneous and mucosal Leishmaniasis 2018

Incidence
(100,000 pop.) **54.7**

Transmission
Triennium Leishmaniasis
Composite Indicator
(TLCIc 2016-2018) **High**

Occurrence


3127

NEW CASES

2673 cutaneous (85.5%)

428 mucosal (13.7%)

26 unspecified (0.8%)



**TLCIc triennium
(2016-2018)**

3.88 - 7.75 (Very intense)

2.65 - 3.83 (Intense)

0.65 - 2.64 (High)

-0.43 - 0.64 (Moderate)

-0.87 - -0.44 (Low)



AGE GROUP

4.4% < 5 Y/O
7.6% ≥ 5 < 10 Y/O
87.7% > 10 Y/O
0.3% Unspecified

12% children under
10 years old



GENDER

70.5% Male
29.5% Female

64% men over 10
years old



CROSS-BORDER

Shares borders with
05 countries in **31**
provinces

36.9% of the cases
at the borders



SURVEILLANCE AND CONTROL

50.5% of the population
in transmission areas

Decrease of the CL % in
children under 10 Y/O
compared to the previous
year (Targets of the 2017-
2022 Plan of Action)



LEISHMANIA SPECIES

Information
NOT available at
SisLeish



LUTZOMYIA VECTORS

Information
NOT available at
SisLeish



PATIENT ASSISTANCE

Laboratory confirmation:
99.9%

Cure rate: **66.8%**

% of cured patients among
the treated: **N/I**



AVAILABLE MEDICINE

Information
NOT available
at SisLeish

Source:

SisLeish - PAHO/WHO - Data provided by the Ministries of Health -
National Leishmaniasis Programs

Accessed on: 01 November 2019.

N/I: No information available; Y/O: years old