

HONDURAS

Leishmaniasis cutaneous and mucosal 2016

Incidence
(100,000 pop.) **43.91**

Transmission
Leishmaniasis
Composite Indicator
(LCIc) **Intense**


Occurrence

2666


NEW CASES

2647 cutaneous (99.3%)

19 mucosal (0.7%)



LCIc



AGE GROUP

10.1% < 5 Y/O
10.7% ≥ 5 < 10 Y/O
79.2% > 10 Y/O

20.8% of children
under 10 years old



GENDER

60.3% Male
39.7% Female

49.8% of men
over 10 years old



CROSS-BORDER

Shares borders with
3 countries in **12**
municipalities

15.1% of the cases
at the borders



SURVEILLANCE AND CONTROL

69.9% of the population
in transmission areas

PEED Participant
(Regional External
Performance Assessment
Program of Microscopic
Diagnosis)



LEISHMANIA SPECIES

L. braziliensis
L. panamensis
L. intantum



LUTZOMYIA VECTORS

L. trapidoi
L. panamensis
L. ylephiletor
L. ovallesi
L. longipalpis



PATIENT ASSISTANCE

Laboratory confirmation:
100% (**24** laboratories
providing diagnosis)

Cure rate: **100%** (**6**
health care units providing
treatment)



AVAILABLE MEDICINE

Meglumine
antimoniate

Deoxycholate
Amphotericin B