

PARAGUAY

Cutaneous and mucosal Leishmaniasis 2017

Incidence (100,000 pop.) **3.73**

Transmission
Triennium Leishmaniasis Composite Indicator (TLCIc 2015-2017) **Low**

Occurrence


92

NEW CASES

30 cutaneous (32.6%)

62 mucosal (67.4%)

LCIc triennium (2015-2017)



AGE GROUP

0% < 5 Y/O
0% ≥ 5 < 10 Y/O
100% > 10 Y/O

0% children under 10 years old



GENDER

79.3% Male
20.7% Female

79.3% men over 10 years old



CROSS-BORDER

Shares borders with **3** countries in **22** municipalities

30.4% of the cases at the borders



SURVEILLANCE AND CONTROL

38.4% of the population in transmission areas

Decrease of the % of CL in children under 10 Y/O compared to the 2012-2015 baseline (Targets of the Plan of Action)



LEISHMANIA SPECIES

L. braziliensis



LUTZOMYIA VECTORS

L. whitmani
L. neivai
L. migonei
L. intermedia



PATIENT ASSISTANCE

Laboratory confirmation: **100%**

Cure rate: **83.7**

% of cured patients among the treated: **N/I**



AVAILABLE MEDICINE

Meglumine Antimoniate

Deoxycholate Amphotericin B

Liposomal Amphotericin B

Source:

SisLeish - PAHO/WHO - Data provided by the Ministries of Health - National Leishmaniasis Programs
Accessed on: 01 November 2018.

N/O: No information available; Y/O: years old